
Purified Rabbit Anti-rat MDC

Catalog Number: TP238

Lot Number: 110819

Content: Protein A purified rabbit IgG, 200 µg, with 0.1% sodium azide, lyophilized.



Product Description and Usage: For research use only. This polyclonal antibody, which reacts with rat MDC, was generated using *E. coli*-expressed rat MDC as an immunogen. The tested titer for Western blot is 1:2,000.

Cross-reactivity to MDC of other species has not been determined.

Storage Condition: 4°C for short term storage or -20°C in small aliquots for long term storage. Avoid repeated freeze and thaw.

Background: MDC (macrophagederived chemokine) is a 69-amino acid CC chemokine with an NH₂-terminal sequence of 24 residues as a signal sequence. It was originally cloned from human monocyte-derived macrophages¹. Recombinant MDC attracts monocytederived dendritic cells. activated lymphocytes and natural killer cells^{1,2}. activity². MDC has anti-HIV-1 Expression of MDC mRNA in

Torrey Pines Biolabs Inc 360 Glenwood Ave, 2nd Fl, Bldg 7 East Orange, NJ 07017 web:www.chemokine.com macrophages is enhanced by LPS, IL-1 β and TNF α^3 . MDC is a functional ligand for CCR4⁴.

References:

- 1. Godiska, R. et al (1997) Human macrophage-derived chemokine (MDC), a novel chemoattractant for monocytes, monocyte-derived dendritic cells, and natural killer cells. *J Exp Med* 185:1595-1604
- 2. Pal, R. et al. (1997) Inhibition of HIV-1 infection by the beta-chemokine MDC. *Science* 278:695-698
- 3. Rodenburg, R.J. (1998) Expression of macrophage-derived chemokine (MDC) mRNA in macrophages is enhanced by interleukin-1beta, tumor necrosis factor alpha, and lipopolysaccharide. *J Leukoc Biol* 63:606-611
- 4. Imai, T. et al. (1998) Macrophage-derived chemokine is a functional ligand for the CC chemokine receptor 4. *J Biol Chem* 273: 1764-1768

Tel: (973)266-8977 Toll free: (866)383-8144 Fax: (973)266-9983 E-mail:info@chemokine.com

